

Influence of Correlated Noises on Growth of a Tumor

Wei-Rong Zhong, Yuan-Zhi Shao,* and Zhen-Hui He

Department of Physics, Sun Yat-sen University, 510275 Guangzhou, People's Republic of China

We studied the effect of additive and multiplicative noises on the growth of a tumor based on a logistic growth model. The steady-state probability distribution and the average population of the tumor cells were given to explain the important roles of correlated noises in the tumor growth. We found that multiplicative noise induces a phase transition of the tumor growth from an unstable state to a bi-stable state; the relationship between the intensity of multiplicative noise and the population of the tumor cells showed a stochastic resonance-like characteristic. It was also confirmed that additive noise weakened rather than extinguish the tumor growth. Homologous noises, however, promote the growth of a tumor. We also discussed about the relationship between the tumor treatment and the model.

PACS numbers: 87.10.+e 05.40.-a 02.50.Ey

I. INTRODUCTION

Nonlinear Physics has changed people's views of world drastically in the past half a century [1, 2], and it is expected useful for discovering the complexity of Biology [3]. More and more attempts confirmed the important role of noise in the nonlinear dynamic systems [4, 5]. Especially, Fokker-Planck equation became one of the approaches of nonlinear dynamics based on deterministic equation with the stochastic form [6, 7]. During the last twenty years, bistable-state model is one of those that attract Physicists most [8-15].

For many years, exploring the growth law of tumors is a challenging subject. Mathematical models are always used as a tentative for describing the tumor growth [16-26]. Bru et al [23, 24] discovered that tumor growth was largely governed by environmental pressures (host tissue pressure, immune response, etc.) through both simulations and experiments of the immune responses of mice. Chemotherapeutic double-faces effect on the cancer was studied by Ferreira et al [25]. It was confirmed that the environment like chemotherapy affected the growth of cancer vitally. Recently, Ai et al [26] applied Logistic growth model to analyze the growth of tumors driven by correlated noises. The relationship between correlated noises and tumor growth was mentioned for the first time. For simplicity, Ai et al only investigated the influence of multiplicative noise on the growth rate of tumors. They found that additive and multiplicative noises would lead the tumors to extinction. They, however, only performed a qualitative analysis according to the steady-state probability distribution of tumor cells. It is unclear whether the influence of correlated noises on a tumor is always negative.

Tumor growth is a complex process. The search based on tumor growth dynamics for the underlying mechanisms of the tumor development and progression might focus on lots of factors. On one hand, the environmental

factors, like temperature, chemotherapy and radiotherapy, may affect the decay rate as well as the growth rate; on the other hand, intrinsic elements, like emigration and genetic instability, also will influence the growth of tumor cells strongly [27]. The environmental factors were regarded as the sources of multiplicative noise, while the intrinsic elements as the additive ones. Considering the fluctuation of the linear growth rate under a stable carrying capacity of environment, we studied the influence of the intensities of correlated noises on the growth and decay of tumor cells. Stochastic resonance (SR) is another concern of this paper. Generally, bistable state is one of the conditions for SR [4]. However, the Logistic model, a non-bistable state system driven by correlated noises as showed in this paper, also shows a typical SR-like characteristic. It is of interest to study this phenomenon as well as its underlying mechanism. We attempt to bridge the gap between correlated noises and tumor treatment and give a new, helpful concept for the treatment of cancer.

II. LANGEVIN EQUATION AND FOKKER-PLANCK EQUATION

Verhulst [28] proposed a single-species model with a self-limiting process. He suggested

$$\frac{dx}{dt} = rx(1 - \frac{x}{K}) \quad (1)$$

This is the Logistic growth equation, where x is the population; r and K are the linear per capita birth rate and the carrying capacity of the environment, respectively. Eq.(1) was used to study the growth of the cells. When the fluctuations of the environment, such as radiotherapy and chemotherapy, were considered, the linear internal growth rate of tumor cell, r , would not be a constant any more. Therefore, it is reasonable to introduce noise $\xi(t)$ into the single-species model, and r should be rewritten as $r + \xi(t)$. Likewise, intrinsic factors were set as another noise $\eta(t)$, e.g. emigration, selection pressure and genetic instability, which could affect the population of tumor cells in local area. So if x was defined as the

*Corresponding Author; Electronic address: stssyz@zsu.edu.cn

population of tumor cells, the Langevin differential equation driven by correlated noises could be written as

$$\frac{dx}{dt} = rx(1 - \frac{x}{K}) + x(1 - \frac{x}{K})\xi(t) + \eta(t) \quad (2)$$

where $\xi(t)$, $\eta(t)$ are respectively the multiplicative and the additive Gaussian white noises defined as

$$\langle \xi(t) \rangle = 0, \quad \langle \xi(t)\xi(t') \rangle = 2D\delta(t - t') \quad (3a)$$

$$\langle \eta(t) \rangle = 0, \quad \langle \eta(t)\eta(t') \rangle = 2Q\delta(t - t') \quad (3b)$$

$$\langle \xi(t)\eta(t') \rangle = 2\lambda\sqrt{DQ}\delta(t - t') \quad (3c)$$

in which D and Q are the intensities of the noises, and λ is the strength of the correlation between $\xi(t)$ and $\eta(t)$ respectively, and $\delta(t - t')$ is Dirac delta function in different times.

According to the Langevin equation (2), one would write out the equivalent Fokker-Planck equation [29].

$$\frac{\partial p(x, t)}{\partial t} = -\frac{\partial}{\partial x}[A(x)p(x, t)] + \frac{\partial^2}{\partial x^2}[B(x)p(x, t)] \quad (4)$$

where $p(x, t)$ is the probability distribution function, $A(x)$ and $B(x)$ are respectively defined as

$$A(x) = rx(1 - \frac{x}{K}) + Dx(1 - \frac{x}{K})(1 - \frac{2x}{K}) + \lambda\sqrt{DQ}(1 - \frac{2x}{K}) \quad (5a)$$

$$B(x) = Dx^2(1 - \frac{x}{K})^2 + 2\lambda\sqrt{DQ}x(1 - \frac{x}{K}) + Q \quad (5b)$$

The population of the cells, x , is positive. According to the reflecting boundary condition, the steady-state probability distribution (STPD) of Eq.(4) is [9]

$$p_{st}(x) = \frac{N}{B(x)} \exp[\int^x \frac{A(x')}{B(x')} dx'] \quad (6)$$

where N is a normalization constant. From this equation, we attained the STPD of the cells population. The peak position of the STPD is K when λ equals zero.

The following parameters are defined to analyze the decay and growth of tumor cells quantitatively.

$$\langle x \rangle = \frac{\int_0^\infty x p_{st}(x) dx}{\int_0^\infty p_{st}(x) dx} \quad (7a)$$

$$\langle x^2 \rangle = \frac{\int_0^\infty x^2 p_{st}(x) dx}{\int_0^\infty p_{st}(x) dx} \quad (7b)$$

$$\langle \delta x^2 \rangle = \langle x^2 \rangle - \langle x \rangle^2 \quad (7c)$$

in which $\langle x \rangle$ and $\langle \delta x^2 \rangle$ refer to the average population of tumor cells and its square difference, respectively. Increasing $\langle x \rangle$ means the growing of the tumor cells, otherwise the decaying. Augment of $\langle \delta x^2 \rangle$ denotes that the tumor growth is unstable, otherwise it is stable.

FIG. 1: The dependence of the STPD on the tumor cells population, the parameters of noises are $\lambda = 0.0$, and $D = 0.5$ respectively. Inset: the average population varies with the intensity of additive noise, from top to bottom, D is 0.20, 0.08 and 0.02, respectively.

III. ANALYSIS OF THE GROWTH MODEL OF A TUMOR

Whether a tumor come into being depends on the linear growth rate, r , in Eq.(1). If r and K are positive, e.g. 0.2 and 6 respectively, the tumors will grow; else if r and K are negative, e.g. -0.2, -6, the tumors will decay. The noises induce different responses of tumor cells between the growth and decay case.

A. Growth Case

The influence of additive noise on the growth of a tumor is shown in Fig.1. In the absence of noises, $p_{st}(x) = \delta(x - K)$, tumors population is stable at $x = K$. In the presence of noises, a STPD appears with its peak at the position $x = K$ (see Fig.1). As the intensity of the noise increases, the peak of STPD decreases. When $D < 0.2$, the population of tumor cells decreases firstly and then increases with the intensity of additive noise, showing a negative stochastic resonance-like characteristic. When $D > 0.2$, the stochastic resonance-like characteristic extincts (see Inset in Fig.1).

In addition to that of the additive noise, the influence of noises correlation on the tumor growth was also studied. The STPD, as shown in Fig.2, increases with the correlation parameter, λ . This implies that the homologous noises promote the stable growth of tumor cells. Inset shows increasing λ is of benefit to tumor growth in the case of low noise intensity; contrarily, increasing λ is unfavorable to the survival of tumor cells. λ has little effect on the tumors at intermediate intensity.

Unlike the additive noise, multiplicative noise has an interesting influence on tumor growth. Fig.3 shows a STPD transition from an uni-peak to a bi-peak distribution with the noise intensity. Moreover, the stochastic resonance

FIG. 2: STPD under different correlations of noises, the noise parameters are $D = 0.5$ and $Q = 2.0$. Inset: the average population varies with the strength of correlation.

FIG. 3: STPD under different multiplicative noise intensities, the parameters are $\lambda = 0.0$ and $Q = 2.0$.

induced by multiplicative noise is also given. In Fig.4, $\langle x \rangle$ increases firstly and then decreases with the intensity of multiplicative noise, showing a typical stochastic resonance characteristic. This means an appropriate intensity of multiplicative noise is suitable for tumor cells, and extra noise restrains tumor growth, even leads them to extinction. Fig.5 shows the influence of noises correlation on the stochastic resonance induced by multiplicative noise. When $\lambda > 0.8$, the stochastic resonance peak will degenerate and disappear. Inset of Fig.5 shows the position D_{peak} , the peak of $\langle x \rangle$, increases exponentially with the parameter, λ .

B. Decay Case

In the decay case, the influence of noise intensity and correlation on the tumors growth is shown in Fig.6-8. Similar to the growth case, in the absence of noises, $p_{st}(x) = \delta(x - 0)$, tumors could not grow. After intro-

FIG. 4: The change of the average population of cells against the intensity of multiplicative noise at $\lambda = 0.0$, from bottom to top, $Q = 0.5, 1.0, 2.0$, respectively.

FIG. 5: The change of the average population of cells with the intensity of multiplicative noise at $Q = 2.0$. Inset: the position, D_{peak} , of the peak of the STPD changes with λ .

ducing noises, one could find that some tumor cells are still alive. In Fig.6, the peak of STPD drops and $\langle \delta x^2 \rangle$ increases with the additive noise intensity. This is in agreement with the results of Fig.1 that additive noise always causes an unstable growth of tumor cells. Different from that of the growth case, the noises correlation has little effect on the STPD of tumor growth in the decay case (see Fig.7). However, inset of Fig.7 shows that the population of tumor cells still increase with the strength of the noises correlation. This denotes that strong correlation would keep the tumor far from extinction. In addition, just like that in the growth case, multiplicative noise also induces a stochastic resonance in the decay case. Fig.8 shows that the peak of STPD increases firstly and then

FIG. 6: The STPD of tumor cells population changes with the intensity of additive noise at $\lambda = 0.0, D = 0.5$. Inset: the varieties of $\langle \delta x^2 \rangle$ with Q .

FIG. 7: STPD under different correlation at $Q = 1.6$ and $D = 0.5$. Inset: the varieties of $\langle x \rangle$ and $\langle \delta x^2 \rangle$ with λ .

decrease with the intensity of multiplicative noise. There is shown in the inset that the average population of cells, $\langle x \rangle$, and the peak of STPD at $x = 0$, $p_{st}(0)$, change with D . It is also a stochastic resonance-like characteristic in the decay case (see the inset of Fig.8).

IV. DISCUSSIONS

Tumor growth was largely governed by their intrinsic elements and the environmental factors like tumor treatment, which was supported through simulations or experiments [24, 25, 26, 30]. In Ref [25], two kinds of selection-inducing therapeutic strategies were considered, namely, those that kill tumor cells and those that block cell mitosis but allowed the cell to survive for some time. These therapeutic strategies were regarded as the external multiplicative noise in our model.

There should be a close relationship between the noises and tumor treatment. Firstly, tumor treatments could

FIG. 8: STPD under different multiplicative noise intensities at $\lambda = 0.0, Q = 1.6$. Inset: the varieties of $\langle x \rangle$ and $p_{st}(0)$ with D .

cause the fluctuation of the tumor cells population. The intensity of multiplicative noise is analogous to the treatment, e.g. the medicinal dose in chemotherapy and the intensity of the ray in radiotherapy. Secondly, the tumor cells themselves transferred from one place to the other ceaselessly to survive or encroach. This caused another fluctuation of the tumor cells. This noise roots in the genetic instability of the tumor cells themselves, so it is internal noise, i.e. additive noise. The intensity of additive noise is considered as an instability of tumor cells population, e.g. genetic instability and cancer heterogeneity. Thirdly, the correlation between two noises reflected the adaptation of a tumor to the treatments. If $\lambda = 0.0$, the tumor was completely unconformable, and if $\lambda = 1.0$, the tumor had better adjustability. Our model did not refer to a certain therapy, but emphasized particularly on the response of a tumor to the treatment.

Additive noise induces an unstable tumor growth, but does not cause extinction. This perhaps owes to the genetic instability and emigration of tumor cells, which adjust themselves to the environment. Furthermore, Multiplicative noise induces a phase transition and stochastic resonance. Firstly, multiplicative noise leads the tumor growth from an uni-stable state to a bi-stable state. This transition denoted the tumor try to survive at the price of partial death. In addition, it was confirmed that tumors grow firstly and then decrease with the enhancement of the treatment. This stochastic resonance-like characteristic implied an improper therapeutic intension could not kill the tumors, but lead to their malignant growth. It was suggested that an enough-strong intensity of therapy was necessary for tumor treatment. Unfortunately, the normal cells were at the risk of canceration at the same time. These presumptions implicated a dilemma for some kinds of therapies like radiotherapy and chemotherapy. Therefore, provided that some advanced treatments based on repeated applications was designed, better treatments would be expected.

It was well known that almost all solid cancers were composed of various genetically and epigenetically more or less different, clonal cell sub-populations. This cancer heterogeneity referred to the fluctuation of tumor cells population as well as a better adaptation of the tumor to the therapies, namely, the additive noise and the correlation of noises. This dual factors focused on an extreme condition, i.e. strong noise with close correlation. Tumor cells population, however, increases steadily under this extreme condition (see inset of Fig.2). This implicated cancer heterogeneity was disadvantageous to cancer curing. It is one of the challenges in modern anti-cancer therapy to apply therapeutic strategies to keep up with the tumor's intrinsic capability to adapt.

The decay case was described as tumor's decreasing under immune surveillance, which caused a tumor to decay. In this case, additive noise still played a role of destroyer like in the growth case, and kept tumor cells far from extinction. Weak multiplicative noise may stabilize the tumor growth, but strong multiplicative noise accelerated their extinctions. However, the noises correlation had little effect on the tumor growth law. These meant that strong-enough treatment and the attempts of different therapies were still necessary for cancer curing even if there existed immune systems.

Anyway, tumor treatment includes kinds of complex processes, such as therapy, immune response, medicine, and so on. Pure therapy just like chemotherapy or radiotherapy has both negative and positive effects on the

tumors. It was suggested in that more measures should be taken in tumor treatment to eliminate a tumor completely, such as gene-therapy, immunotherapy, resection and so on.

V. CONCLUSIONS

We presumed two kinds of noises in the tumor growth model. One originates from the tumor treatment, like radiotherapy and chemotherapy; the other roots in the tumor cells themselves. The previous is multiplicative, and the latter is additive. On one hand, multiplicative noises induce a phase transition and stochastic resonance in the tumor growth. An appropriate intensity of multiplicative noise, e.g. an error treatment, could not kill a tumor, but activated it. On the other hand, additive noises cause an unstable growth of a tumor, but does not extinct it. Under a weak multiplicative noise, an additive noise also induces a negative stochastic resonance. In addition, close correlation between multiplicative and additive noises, i.e. a better adjustability of a tumor to the environment, is benefit to the growth of tumors.

ACKNOWLEDGEMENTS

This work was partially supported by the National Natural Science Foundation (Grant No. 60471023) and the Natural Science Foundation of Guangdong Province (Grant No. 031554), P. R. China.

-
- [1] G. Nicolis and I. Prigogine, *Self-organization in Nonequilibrium systems* (Wiley New-York, 1977).
 - [2] H. Haken, *Advanced Synergetics* (Springer Berlin Heidelberg, 1985).
 - [3] G. Nicolis and I. Prigogine, *Exploring Complex* (Freeman New-York, 1986).
 - [4] L. Gammaitoni, P. Hanggi, P. Jung, and F. Marchesoni, *Rev. Mod. Phys.* 70, 223 (1998).
 - [5] V. S. Anishchenko, V. V. Astakhov, A. B. Neiman, T. E. Vadivasova, and L. Schimansky-Geier, *Nonlinear Dynamics of Chaotic and Stochastic Systems* (Springer-Verlag Berlin Heidelberg, 2002).
 - [6] C. W. Gardiner, *Handbook of Stochastic Methods for Physics, Chemistry and the Natural Science* (Springer-Verlag Berlin, 1983).
 - [7] G. Hu, *Stochastic Forces and Nonlinear Systems* (Shanghai Scientific and Technological Education Publishing House, Shanghai, P. R. China, 1994) (in Chinese).
 - [8] Y. Jia and J. R. Li, *Phys. Rev. Lett.* 78, 994 (1997).
 - [9] Y. Jia and J. R. Li, *Phys. Rev. E* 53, 5786 (1996).
 - [10] P. Hanggi and P. Riseborough, *Phys. Rev. A* 27, R3379 (1983).
 - [11] B. McNamara, K. Weisenfeld and R. Roy, *Phys. Rev. Lett.* 60, 2626 (1988).
 - [12] G. Hu, G. Nicolis, and C. Nicolis, *Phys. Rev. A* 42, 2030 (1990).
 - [13] P. Jung and P. Hanggi, *Phys. Rev. A* 44, 8032 (1991).
 - [14] M. I. Dykman, R. Mannella, P. V. E. McClintock, and N. G. Stocks, *Phys. Rev. Lett.* 68, 2985 (1992).
 - [15] A. A. Zaikin, J. Kurths, and L. Schimansky-Geier, *Phys. Rev. Lett.* 85, 227 (2000).
 - [16] J. D. Murray, *Mathematical Biology I: An Introduction* (Springer-Verlag Berlin Heidelberg, 2002).
 - [17] R. B. Banks, *Growth and Diffusion Phenomena* (Springer-Verlag Berlin Heidelberg, 1994).
 - [18] M. Scalerandi and F. Peggion, *Phys. Rev. E* 66, 031903 (2002).
 - [19] J. Yu, G. Hu, and M. B. Kun, *Phys. Rev. B* 39, 4572 (1989).
 - [20] M. Molski and J. Konarski, *Phys. Rev. E* 68, 021916 (2003).
 - [21] S. Kar, S. K. Banik, and D. S. Ray, *Phys. Rev. E* 65, 061909 (2002).
 - [22] M. Scalerandi and B. C. Sansone, *Phys. Rev. Lett.* 89, 218101 (2002).
 - [23] A. Bru, J. M. Pastor, I. Fernaund, I. Bru, S. Melle, and C. Berenguer, *Phys. Rev. Lett.* 81, 4008 (1998).
 - [24] A. Bru, S. Albertos, J. A. Lopez. Garcia-Asenjo, and I. Bru, *Phys. Rev. Lett.* 92, 238101 (2004).
 - [25] S. C. Ferreira, Jr, M. L. Martins, and M. J. Vilela, *Phys. Rev. E* 67, 051914 (2003).
 - [26] B. Q. Ai, X. J. Wang, G. T. Liu, and L. G. Liu, *Phys. Rev. E* 67, 022903 (2003).
 - [27] R. V. Sole and T. S. Deisboeck, *J. Theor. Biol.* 228, 47 (2004).
 - [28] P. -F. Verhulst. *Recherche mathématiques sur le loi*

- d'accroissement de la population. Nouveau Memoirès de l'Académie Royale des Sciences et Belles Lettres de Bruxelles, 18:3-38, (1845).
- [29] W. D. Jin, C. Li, and K. S. Zhi, Phys. Rev. E 50, 2496 (1994).
- [30] A. Bru, S. Albertos, J. L. Subiza, J. L. Garcia-Asenjo, and I. Bru, Biophys. J. 85, 2948 (2003).















